

Emergency Bodily Fluid Clean-Up [Ready-to-Use]

Products Needed



Comet®
Disinfecting
Bathroom
Cleaner

Tools Needed

- 2 Garbage Bags with Ties
- Baking Soda (optional)
- Disposable PPE (see Step 2 below)
- Paper Towels
- Safety Cones/Signs

Frequency


As Needed


Special Directions

Cleanup of blood, vomit, diarrhea, and other bodily excretions should be handled in an urgent manner. Staff should be trained in proper use of personal protective equipment.

1  Remove all individuals from affected area (up to a 25 foot radius).

2 Gather supplies including disposable personal protective equipment (PPE): 1) goggles; 2) face mask 3) gloves; 4) shoe covers; 5) apron or gown.

3  Put on PPE: 1) goggles; 2) face mask; 3) gown or apron; 4) gloves; and 5) shoe covers.

4  Sprinkle baking soda powder over bodily fluid incident and/or cover with paper towels to absorb.

5  Spray disinfecting cleaner over the affected area and about 3 feet beyond.

NOTE: Carpet and soft furnishings should be cleaned with steam at 70 °C (158 °F) or higher for 5 minutes.


6  Dispose of exposed food within a 25 foot radius of event, if applicable.


7  Wipe up solidified soils with towels and then throw them away in the trash bag.


8  Spray, thoroughly, with disinfecting surface cleaner over affected area and 3 feet beyond and allow a 10 minute contact time.

9  Wipe up cleaner and soil residues with clean, dry towels and throw away in trash bag.

10  Spray disinfecting cleaner on high touch surfaces (table tops, chairs, door knobs, etc) within a 25 foot radius of vomitus event and allow a 10 minute contact time, if applicable.



11  Tie first bag and place it in the second trash bag. Remove PPE and throw away in second trash bag. Remove gloves last and throw away in second trash bag. Dispose of biohazardous waste in accordance with local regulations.

12  Wash hands, thoroughly.

every experience counts.™

P&G
Professional